

# Exhibit 13

David A. Bahr (D.D.C. Bar # OR0001)  
1035 ½ Monroe St  
Eugene, OR 97402  
(541) 566-6439  
[davebahr@mindspring.com](mailto:davebahr@mindspring.com)

Kristine Akland (Montana Bar # 13787) (*pro hac vice*)  
317 E Spruce St, P.O. Box 7274  
Missoula, MT 59807  
(406) 544-9863  
[aklandlawfirm@gmail.com](mailto:aklandlawfirm@gmail.com)

John Persell (D.D.C. Bar # ID0002)  
Western Watersheds Project  
P.O. Box 1770  
Hailey, ID 83333  
(503) 896-6472  
[jpersell@westernwatersheds.org](mailto:jpersell@westernwatersheds.org)

*Attorneys for Plaintiffs*

**IN THE UNITED STATES DISTRICT COURT  
FOR THE DISTRICT OF COLUMBIA**

WESTERN WATERSHEDS PROJECT,  
ALLIANCE FOR THE WILD  
ROCKIES, and YELLOWSTONE TO  
UINTAS CONNECTION

Plaintiffs,

v.

DAVID BERNHARDT, in his official  
capacity as Secretary, U.S. Department  
of the Interior, UNITED STATES FISH  
AND WILDLIFE SERVICE, and  
UNITED STATES FOREST SERVICE,

Defendants.

**Civil Action No. 1:20-cv-860-APM**

**DECLARATION OF DR. DAVID J. MATTSON**

I, DAVID J. MATTSON, hereby declare:

1. I am a scientist and retired wildlife management professional with extensive experience in grizzly bear research and conservation spanning four decades. My educational attainments include a B.S. in Forest Resource Management, an M.S. in Plant Ecology, and a Ph.D. in Wildlife Resource Management. My professional positions prior to retirement from the U.S. Geological Survey (USGS) in 2013 included: Research Wildlife Biologist, Leader of the Colorado Plateau Research Station, and Acting Center Director for the Southwest Biological Science Center, all with the USGS; Western Field Director of the Massachusetts Institute of Technology-USGS Science Impact Collaborative; Visiting Scholar at the Massachusetts Institute of Technology; and Lecturer and Visiting Senior Scientist at the Yale School of Forestry & Environmental Studies.

My dissertation focused on the ecology of grizzly bears in the Greater Yellowstone Ecosystem (GYE) during 1977-1996 (Mattson 2000). I intensively studied grizzly bears in the GYE during 1979-1993 as part of the Interagency Grizzly Bear Study Team (IGBST) and was charged with designing and supervising field investigations during 1985-1993. My field research focused on human-grizzly bear relations; grizzly bear foraging, habitat selection, diet, and energetics; and availability and ecology of grizzly bear foods. I have continued to closely observe grizzly bears and their habitats in the GYE since the end of my intensive field investigations in 1993.

Although my field studies in the GYE ended in 1993, my involvement in grizzly bear-related research, management, and education, both regionally and internationally, has continued through the present. Throughout my career I have

been consulted by brown/grizzly bear managers and researchers worldwide, including from Russia, Japan, France, Spain, Greece, Italy, and, most notably, Canada. I have also given numerous public presentations on grizzly bear ecology and conservation, including talks, nationally, at the Smithsonian (Washington, DC) and American Museum of Natural History (New York, NY), and, regionally, at the Denver Museum of Natural History (Denver, CO), the Museum of Wildlife Art (Jackson, WY), and the Museum of the Rockies (Bozeman, MT).

2. What follows is my assessment of the analysis of impacts on grizzly bears reported in the Environmental Impact Statement (EIS) and Biological Opinion (BiOP) for grazing operations on US Forest Service allotments in the Upper Green River (UG) area of the Bridger-Teton National Forest. I start with some necessary ecological context, then address relevant temporal and geospatial dynamics in the Greater Yellowstone Ecosystem (GYE), and end with my evaluation of the analyses and related conclusions reported in the EIS and BiOP.

Attachment 1 contains the scientific literature and sources that I cite in this declaration. Attachment 2 contains figures and figure captions that I reference throughout.

## ECOLOGICAL CONTEXT

3. Meat from livestock and other large herbivores is a high-quality grizzly bear food (Mattson et al. 2004; Erlenbach et al. 2014).

4. Male bears tend to eat more meat. However, if meat is one of the few high-quality foods available to bears, levels of meat consumption by male and female bears will converge (Jacoby et al. 1999; Mattson 1997, 2000; Hobson et al. 2000; McLellan 2011; Mace & Roberts 2012; Fortin et al. 2013; Schwartz et al.

2013).

5. Some level of depredation by bears will predictably occur if livestock are available and vulnerable (Murie 1948, Johnson & Griffell 1982, Bjorge 1983, Knight & Judd 1983, Jorgensen 1983).

6. Availability of vulnerable livestock or human-associated carrion can attract bears to an area and lead to local increases in bear densities (Bailey 1931, Brown 1996, Storer & Tevis 1996, Mattson & Merrill 2002, Haroldson et al. 2004).

7. Grizzly bears eat more meat from large herbivores when other high-quality foods are not as abundant; bears can also be a significant source of predation on large herbivores (Mattson 1990, 1997; Mowat & Heard 2006; Zager & Beecham 2006; Barber-Meyer et al. 2008; Vulla et al. 2009; Middleton et al. 2013; Schwartz et al. 2013; Ebinger et al. 2016; Niedzialkowska et al. 2018).

8. Dependence of grizzly bears on meat from livestock leads to higher mortality rates, reduced densities and local extirpations under regimes where lethal control is the primary human response (Storer & Tevis 1996, Brown 1996, Mattson & Merrill 2002, Merrill & Mattson 2003, Mowat et al. 2013).

#### CONCLUSIONS: Ecological Context

9. These basic observations lead to some unambiguous prefatory conclusions, including: (i) consumption of meat from livestock by grizzly bears is normal and to be expected given the high quality of this food resource; (ii) some level of predation on livestock is also normal, expected, and even inevitable any time livestock and grizzly bears share space; (iii) consumption of meat from large herbivores will increase as abundance of other high-quality foods lessens, either

inter-annually, as a trend over time, or as a function of intrinsic ecological conditions; and (iv) livestock can be an attractant, a local driver of increased bear densities, as well as more often a catalyst for lethal relations with humans that lead to sometimes dramatic reductions in bear numbers, followed by local extirpations.

THE GREATER YELLOWSTONE ECOSYSTEM (GYE) AS CONTEXT FOR THE UPPER GREEN (UG)

10. Seeds from whitebark pine cones were once a critically important food for GYE grizzly bears, more so for female bears than for male bears (Mattson et al. 1991a, 2004; Mattson 2000).

11. When whitebark pine was still an abundant part of GYE grizzly bear habitat, conflicts with as well as exposure to humans increased during years when whitebark pine seeds were scarce (Mattson et al. 1992, Haroldson & Gunther 2013), resulting in higher levels of bear mortality during years with poor whitebark pine seed crops (Mattson et al. 1992, Mattson 1998, Pease & Mattson 1999, Schwartz et al. 2006).

12. Roughly 70% of mature cone-producing whitebark pine trees were lost in the GYE between 2000 and 2010 to a climate-driven outbreak of mountain pine beetles (Macfarlane et al. 2013; Van Manen et al. 2016, 2019; Figure 1c).

13. Losses of trees were most pronounced in the central and southern Absaroka Mountains and in the UG area (Macfarlane et al. 2013; Figure 1b).

14. Losses of whitebark pine *seeds* accelerated after 2007. Ecosystem-wide, after accounting for both tree losses and production of cones per tree, seed availability has remained at low levels since then (Van Manen et al. 2016; data from: Schwartz & Haroldson 1999-2004; Schwartz et al. 2005-2011; Van Manen

et al. 2012-2019; Figure 1c-e)

15. Consumption of meat from large herbivores by grizzly bears in the GYE has steadily increased since the early 2000s, likely in compensation for losses of whitebark pine and cutthroat trout (Orozco & Miles 2012; Middleton et al. 2013; Schwartz et al. 2013; Ebinger et al. 2015; Figure 2a-b), at the same time that numbers of elk declined along with numbers of ungulate carcasses available for spring foraging by bears on winter ranges (Figure 2c-e).

16. Army cutworm moths are another high-quality bear food intensively used by grizzly bears, primarily in the central Absaroka Mountains. Moth sites are notably absent in and near the UG area (Mattson et al. 1991b, French et al. 1994, Robison 2009; Figure 3b,f).

17. Although numbers of moth sites have not increased, levels of bear activity on these sites have increased substantially since 2010, likely in compensation for losses of whitebark pine (data from: Van Manen et al. 2019; Figure 3c,d).

18. Trend of the reproductive segment of the grizzly bear population within the GYE Demographic Monitoring Area (DMA) has been stable to only slightly increasing since the early 2000s (data from: Van Manen et al. 2019; Figures 5,6).

19. Total known and probable deaths of adolescent and adult grizzly bears within the GYE have been increasing at annual rates greater than any possible increase in total size of the bear population (data from: Schwartz & Haroldson 1999-2004; Schwartz et al. 2005-2011; Van Manen et al. 2012-2019, Figure 6).

20. Most of this increase in bear mortality has been driven by deaths

related to conflicts with humans over meat resources, including livestock and carcasses associated with elk hunters (data from: database for 1959-2014 obtained under terms of Federal FOIA; Schwartz & Haroldson 1999-2004; Schwartz et al. 2005-2011; Van Manen et al. 2012-2019; Figure 2g,h & Figure 4c).

21. In common with other grizzly bear populations, growth of the Yellowstone grizzly bear population is more sensitive to annual survival of adult females than to any other single vital rate (e.g., Knight & Eberhardt 1985, Hovey & McLellan 1996, Schwartz et al. 2006). In other words, increase or decline of the population will be strongly affected by the rates at which reproductive-aged females are recruited and survive. In recognition of this basic fact, current protocols governing management of mortality for the Yellowstone grizzly bear population prescribe an allowable rate of mortality for adolescent and adult female bears that is only half that allowed for adolescent and adult males (Wyoming Game & Fish Department 2016).

22. Distribution of the GYE grizzly bear population encompassed the UG area as early as 1998 and, overall, has increased at an annual rate 4- to 26-times greater than increases in population size (data from: Van Manen et al. 2019; Figure 1a,b). Increases in distribution cannot be explained solely by increases in bear numbers.

23. Depredation-related conflicts in the UG area escalated dramatically after 2010 along with lethal removal of depredating grizzly bears (data from: Schwartz & Haroldson 1999-2004; Schwartz et al. 2005-2011; Van Manen et al. 2012-2019; Figures 4d, 5, 7).

24. Female grizzly bears have accounted for a significant portion of



depredation-related removals after but not before 2010 (data from: Schwartz & Haroldson 1999-2004; Schwartz et al. 2005-2011; Van Manen et al. 2012-2019; Figure 7).

25. Concentrations of depredations and depredation-related conflicts are 9- to 24-times greater in the UG grazing allotments (depending on time period) compared to any other single allotment or complex of allotments in the GYE (data from: Schwartz & Haroldson 1999-2004; Schwartz et al. 2005-2011; Van Manen et al. 2012-2019)—in defiance of any straight-forward relationship with local landscape features (Figure 8).

26. Escalation of depredations in the UG after 2010 is not temporally or spatially correlated with increases in size and distribution of the GYE grizzly bear population (data from: Schwartz & Haroldson 1999-2004; Schwartz et al. 2005-2011; Van Manen et al. 2012-2019; Figure 9).

27. At a broad scale, escalation of depredations in the UG is strongly positively correlated, temporally and spatially, with losses of whitebark pine and absence of army cutworm moth sites (Mattson et al. 1991b, Macfarlane et al. 2013, Robison 2009; Figures 4b,d and Figure 9).

28. At a finer scale, level of depredation is correlated with the existence of remnant stands of whitebark pine that presumably continue to attract grizzly bears into areas near the UG (Costello et al. 2014, Wells et al. 2019).

29. Dynamics associated with depredation of livestock in the UG strongly resemble those of an “ecological trap” such as has been described for grizzly and brown bears elsewhere (Knight et al. 1988, Falcucci et al. 2009, Northrup et al. 2012, Lamb et al. 2017).

## CONCLUSIONS: The GYE as context for the UG

30. Weight of evidence overwhelming supports concluding that: (i) livestock on grazing allotments in the UG serve as an attractive high-quality food for most grizzly bears in the area; (ii) grizzly bears in this area have turned increasingly to exploiting livestock to compensate for losses of whitebark pine, with this diet switch amplified by scarcity of high-quality alternative foods such as cutworm moths and the nearness of remnant stands of whitebark pine to extant grazing allotments; (iii) livestock in the UG fuel an “ecological trap,” attracting bears of both sexes into situations that end up being lethal to the involved bears; and (iv) there is not a surfeit of grizzly bears in the DMA able to sustain the locally elevated rate of mortality associated with lethal response to depredation in the UG.

## GEOSPATIAL CONTEXT

31. Five different scientific analyses have shown there to be ample potential suitable habitat for grizzly bears south and southeast of the UG in the Wind River and Wyoming Ranges (Carroll et al. 2003, Merrill & Mattson 2003, Craighead et al. 2005, Merrill 2005, Schwartz et al. 2010; Figure 10).

32. Expansion of the GYE grizzly bear population since 1990 has confirmed the validity of these prior analyses (Van Manen et al. 2019).

33. The UG is located in a critical area separating core and peripheral grizzly bear habitat in the GYE (Merrill & Mattson 2003, Craighead et al. 2005, Merrill 2005, Schwartz et al. 2010; Figure 10).

34. Roads and livestock-related mortalities associated with UG grazing allotments are associated with a zone of elevated mortality rates separating core and peripheral grizzly bear habitat (Merrill & Mattson 2003, Schwartz et al. 2010;

Figure 10a,b).

## CONCLUSIONS: Geospatial Context

35. The overwhelming weight of evidence supports concluding that: (i) the UG area is important to continuity between core and peripheral habitat grizzly bear habitat in the GYE; (ii) peripheral habitat in the Wind River Range beyond the UG supports a substantial portion of the bears required to meet demographic objectives within the DMA; and (iii) high levels of road access and livestock-related grizzly bear mortality within the UG have created one of the most pronounced fracture zones within the GYE.

## MITIGATION MEASURES

36. There is some evidence suggesting that selective removal of a few depredating grizzly bears (e.g., 1-3 over a period of 1-2 years) can resolve conflicts for short periods of time (e.g., 1-3 years), but contingent on specific favorable circumstances (Anderson et al. 2002, Miller et al. 2016, Morehouse et al. 2016, Swan et al. 2017, Lennox et al. 2018, Proulx 2018).

37. There is little or no evidence that supports the effectiveness or conservation-efficacy of removing numerous bears every year to resolve depredation-related conflicts, especially under circumstances where livestock are a primary high-quality food fueling an “ecological trap” (e.g., Sagør et al. 1997, Graham et al. 2005, Treves & Naughton-Treves 2005, Northrup et al. 2012, Miller et al. 2016, Morehouse et al. 2016, Treves et al. 2016, Eklund et al. 2017, Lamb et al. 2017, Swan et al. 2017, Lennox et al. 2018, Lute et al. 2018, Moreira-Arce et al. 2018, Proulx 2018, Haswell et al. 2019). Rather, as in the UG, most evidence suggests that conflicts continue unabated until bears are locally extirpated.

38. There is both correlative and causational evidence suggesting that non-lethal measures can reduce levels of depredation for sustained periods of time, including: guardian dogs; selective deployment of electric fence and other deterrents; change in species, sex, and age of grazed livestock; closer guarding; relocation of pastures during key periods of livestock vulnerability; and removal of livestock carcasses (e.g., Bjorge 1983; Wilson et al. 2005, 2006; Karlsson & Johansson 2010; Miller et al. 2016; Treves et al. 2016; Eklund et al. 2017; Moreira-Arce et al. 2018; Khorozyan & Waltert 2019a, 2019b; Wells et al. 2019).

39. There is definitive evidence from the GYE showing that retirement of livestock grazing allotments eliminates or precludes depredation-related conflicts with bears (e.g., lack of depredations in retired allotments [Wells 2017, Wells et al. 2019]; also, lack of depredations after retirement on the Blackrock-Spread Creek allotment immediately to the west of the UG where conflicts had been chronic since before the 1930s [Murie 1948, Knight & Judd 1983, Anderson et al. 2002]; and lack of depredations after retirement of sheep grazing allotments with a long history of chronic conflict [Johnson & Griffel 1982, Knight & Judd 1983, Jorgensen 1983, Van Manen et al. 2019]; see Figure 4a).

#### CONCLUSIONS: Mitigation Measures

40. Weight of evidence supports concluding that: (i) short of causing near-term local extirpation of grizzly bears, continued lethal removal of depredating bears will not appreciably resolve conflicts in the UG; (ii) sustained deployment of non-lethal measures stands a good chance of reducing levels of depredation; and (iii), given the evident drivers, the most effective means for resolving depredation-related conflicts in the UG is retirement of allotments

suffering chronic conflict.

## EVALUATION OF THE UPPER GREEN EIS AND BIOP

41. The US Forest Service EIS and related US Fish & Wildlife Service BiOP justify future grazing operations in the UG are based on the following series of explicit or tacit claims: (i) Conflicts/depredations in the UG have increased solely because of an increasing and expanding grizzly bear population; (ii) Prospective non-lethal mitigation measures are unproven, infeasible, or precluded by negative side-effects; (iii) Killing bears is the only feasible way of preventing and resolving conflicts in the UG; (iv) Because the bear population is increasing and expanding and will continue to do so in the future, managers can kill as many bears in the UG during the next 10 years as they have in the past; (v) There is no need to factor future or recent changes in population-wide bear mortality into mortality that is allowed in the UG during the next 10 years; (vi) High levels of mortality in the UG are not problematic at a landscape level because there is little or no potential suitable habitat for grizzlies to the south and southeast; (vii) Depredation-related grizzly bear mortality and conflicts in the UG are not exceptional; and (viii) Female bears are as expendable as male bears.

41. Claims (i), (ii), (iii), (v), (vi), and (vii) contradict or are inconsistent with the best available science, as summarized above. Increased conflicts are not correlated with changes in grizzly bear population size and distribution. Instead, this increase is positively correlated with loss of whitebark pine and a dearth of nearby moth sites that would otherwise accommodate a switch to eating cutworm moths. There is no evidence supporting the efficacy of killing bears to resolve depredation conflicts when conflicts are likely organized around an “ecological

trap.” Instead, there is good evidence supporting the efficacy of various non-lethal strategies, foremost of which is retiring allotments with chronic conflict. And, finally, the bear population within the DMA has been nearly static since conflicts, conflict-related bear deaths, and ecosystem-wide bear deaths have escalated during the last 6 years, which does not support the contention that the population has absorbed and can continue to absorb bear deaths in the UG without some degree of harm.

42. Claim (v) is problematic because of a logical disconnect. The Memorandum-of-Agreement (MOA) guiding current management of the GYE grizzly bear population adjusts allowable or target mortality rates based on estimates of total grizzly bear population size (Wyoming Game & Fish Department 2016). If the estimated population declines, progressively lower rates are applied. As a consequence, fewer bears can be killed population-wide each year if the population is to be sustained. Current arrangements require that consultation with the USFWS be reinitiated only if numbers of bears killed within or near the UG surpass a certain threshold, not if mortality thresholds for the entire population have been lowered or surpassed. This oversight could allow bears to be killed in the UG as a percent of total estimated population size at a rate that is higher in the future than in the past if the DMA population has declined, which contradicts the assumption that future mortality in the UG will take no more of a toll on the population than has past mortality. This risk is not addressed in either the EIS or BiOP.

43. Of relevance to claim (v) as well, the situation in the UG is clearly dynamic. This fact is evident in the extent to which conflicts escalated and average

bear mortality increased after 2010 along with the presence of female bears in this toll. Absent any real-time monitoring of conditions driving these increases, a moving average of bear deaths will be tantamount to managing through a rearview mirror and will, moreover, offer no insight into reasons why changes may have occurred. Without such insight, managers will be handicapped when trying to identify options for rectifying a worsening situation.

44. The short-comings of relying primarily on a running-average of past mortality for management of grizzly bears in the UG are exacerbated by the fact that the Upper Green EIS and BiOP fail to consider foreseeable effects of climate change. The relevant body of science is too large to allow for even a cursory summary here. At the very least, we can expect more extensive and frequent wildfires (e.g., Westerling et al. 2011, Liu & Wimberly 2016, Clark et al. 2017, Gergel et al. 2017), distorted hydrologic regimes (e.g., Littell et al. 2011, Luce 2018), loss of wetlands (Ray et al. 2019), changes in amounts and seasonal availability of forage (e.g., Reeves et al. 2014, Hufkens et al. 2016, Thoma et al. 2019), little or no recovery of whitebark pine and cutthroat trout (e.g., Williams et al. 2009; Isaak et al. 2012; Al-Chokhachy et al. 2013, 2017; Chang et al. 2014; Case & Lawler 2016; Buotte et al. 2016, 2017), reductions in amounts of berries (Prev  y et al. 2020), and collapse of alpine environments (e.g., Rehfeldt et al. 2012, Hansen & Phillips 2015) along with alpine-associated army cutworm moth sites (Mattson et al. 1991b, White et al. 1998). The EIS and BiOP offer no assessment of these foreseeable impacts on grizzly bears and, for unexplained or poorly justified reasons, assume that post hoc monitoring of mortality will be sufficient for addressing this momentous issue.

45. Of relevance to the Government's claim (viii), the UG EIS does not acknowledge the toll that mortality has taken on females grizzly bears since 2010 and also fails to make any provision for the preferential protection of females. As I note under point 21 above, growth of the Yellowstone grizzly bear population is far more sensitive to the rates at which reproductive females are recruited and survive than to the same rates for male bears. Lack of stringent protections for female grizzly bears in the UG EIS is thus both alarming and a fatal short-coming.

CONCLUSIONS: Failings of the UG EIS and BiOP

46. The Upper Green EIS and BiOP are fatally flawed and offer an approach to managing grizzly bears in this region that exacerbates rather than alleviates threats. For one, key issues are not addressed. For another, the reported analyses use only a selection of relevant scientific information, with this selectivity appearing to serve preordained ends. Perhaps more disturbing, grizzly bears of both sexes are assumed to be expendable. As an apparent consequence, the efficacy of lethal management is grossly over-stated and the potential benefits of non-lethal methods given short-shrift. Most important of all, the alternative of terminating grazing on allotments suffering chronic conflict is not given appropriate consideration. This omission is especially troubling given that this option that has been widely implemented elsewhere in the GYE.

As Richard Knight and Steven Judd observed several decades ago, "We believe that cattle allotments can be permitted in grizzly bear habitat if it is understood that the cattle owners are willing to absorb grizzly bear predation losses" (Knight & Judd 1983). This sort of perspective—voiced by respected wildlife researchers—apparently did not guide either the US Forest Service or US



Fish & Wildlife Service. The EIS is neither precautionary nor prudent, but rather profligate with grizzly bears protected by the Endangered Species Act.

47. For all of the reasons articulated above, it is my expert opinion that: (i) analyses pertaining to impacts on grizzly bears presented in the Upper Green EIS and supporting BiOP are patently inadequate and at odds with the best available science; and (ii) grazing operations and grizzly bear management planned under terms of the Upper Green EIS will needlessly harm not only individual grizzly bears and the GYE Yellowstone grizzly bear population, but also long term conservation of this population.

Pursuant to 28 U.S.C. § 1746, I declare under penalty of perjury that the foregoing is true and correct.

Executed on this 7<sup>th</sup> day of May, 2020.

A handwritten signature in black ink, reading "David J. Mattson". The signature is written in a cursive style with a large, looped initial "D".

David J. Mattson, Ph.D.

## **Attachment 1. Literature Cited in Main Text.**

- Al-Chokhachy, R., Alder, J., Hostetler, S., Gresswell, R., & Shepard, B. (2013). Thermal controls of Yellowstone cutthroat trout and invasive fishes under climate change. *Global Change Biology*, 19(10), 3069-3081.
- Al-Chokhachy, R., Sepulveda, A. J., Ray, A. M., Thoma, D. P., & Tercek, M. T. (2017). Evaluating species-specific changes in hydrologic regimes: an iterative approach for salmonids in the Greater Yellowstone Area (USA). *Reviews in Fish Biology & Fisheries*, 27(2), 425-441.
- Anderson Jr, C. R., Terner, M. A., & Moody, D. S. (2002). Grizzly bear-cattle interactions on two grazing allotments in northwest Wyoming. *Ursus*, 13, 247-256.
- Aune, K., & Kasworm, W. (1989). East Front grizzly studies: final report. Montana Department of Fish, Wildlife & Parks, Helena, Montana.
- Bailey, V. (1931). Mammals of New Mexico. *North American Fauna*, 53, 349-368.
- Barber-Meyer, S. M., Mech, L. D., & White, P. J. (2008). Elk calf survival and mortality following wolf restoration to Yellowstone National Park. *Wildlife Monographs*, 169(1), 1-30.
- Bjorge, R. R. (1983). Mortality of cattle on two types of grazing areas in northwestern Alberta Canada, woodland pastures, intensively managed pastures. *Journal of Range Management*, 36(1), 20-21.
- Brown, D. E. (1996). *The Grizzly in the Southwest: Documentary of an Extinction*. University of Oklahoma Press, Norman, Oklahoma.
- Buotte, P. C., Hicke, J. A., Preisler, H. K., Abatzoglou, J. T., Raffa, K. F., & Logan, J. A. (2016). Climate influences on whitebark pine mortality from mountain pine beetle in the Greater Yellowstone Ecosystem. *Ecological Applications*, 26(8), 2507-2524.
- Buotte, P. C., Hicke, J. A., Preisler, H. K., Abatzoglou, J. T., Raffa, K. F., & Logan, J. A. (2017). Recent and future climate suitability for whitebark pine mortality from mountain pine beetles varies across the western US. *Forest Ecology & Management*, 399, 132-142.
- Carroll, C., Noss, R. F., Paquet, P. C., & Schumaker, N. H. (2003). Use of population viability

- analysis and reserve selection algorithms in regional conservation plans. *Ecological Applications*, 13(6), 1773-1789.
- Case, M. J., & Lawler, J. J. (2016). Relative vulnerability to climate change of trees in western North America. *Climatic Change*, 136(2), 367-379.
- Chang, T., Hansen, A. J., & Piekielek, N. (2014). Patterns and variability of projected bioclimatic habitat for *Pinus albicaulis* in the Greater Yellowstone Area. *PLoS One*, 9(11).
- Chapman, J. A., Romer, J. I., & Stark, J. (1955). Ladybird beetles and army cutworm adults as food for grizzly bears in Montana. *Ecology*, 36(1), 156-158.
- Chapman, S.S., Bryce, S.A., Omernik, J.M., Despain, D.G., ZumBerge, J., & Conrad, M. (2004). Ecoregions of Wyoming (color poster with map, descriptive text, summary tables, and photographs): Reston, Virginia, U.S. Geological Survey, Reston, Virginia.
- Clark, J. A., Loehman, R. A., & Keane, R. E. (2017). Climate changes and wildfire alter vegetation of Yellowstone National Park, but forest cover persists. *Ecosphere*, 8(1), e01636.
- Costello, C. M., van Manen, F. T., Haroldson, M. A., Ebinger, M. R., Cain, S. L., Gunther, K. A., & Bjornlie, D. D. (2014). Influence of whitebark pine decline on fall habitat use and movements of grizzly bears in the Greater Yellowstone Ecosystem. *Ecology & Evolution*, 4(10), 2004-2018.
- Craighead, J. J., Sumner J. S., Scaggs, G. B. (1982). A definitive system for analysis of grizzly bear habitat and other wilderness resources. Craighead Western Wildlands Institute Monograph 1. University of Montana Foundation, Missoula, Montana.
- Craighead, L., Gilbert, B., & Olenicki, T. (2005). Comments submitted to the US Fish and Wildlife Service regarding delisting of the Yellowstone Grizzly Bear DPS, Federal Register. Vol. 70, No. 221. (November 17, 2005): 69853–69884.
- Ebinger, M. R., Haroldson, M. A., Van Manen, F. T., Costello, C. M., Bjornlie, D. D., Thompson, D. J., ... & White, P. J. (2016). Detecting grizzly bear use of ungulate carcasses using global positioning system telemetry and activity data. *Oecologia*, 181(3), 695-708.
- Eklund, A., López-Bao, J. V., Tourani, M., Chapron, G., & Frank, J. (2017). Limited evidence on the effectiveness of interventions to reduce livestock predation by large carnivores.

- Scientific Reports, 7(1), 1-9.
- Erlenbach, J. A., Rode, K. D., Raubenheimer, D., & Robbins, C. T. (2014). Macronutrient optimization and energy maximization determine diets of brown bears. *Journal of Mammalogy*, 95(1), 160-168.
- Falcucci, A., Ciucci, P., Maiorana, L., Gentile, L., & Boitani, L. (2009). Assessing habitat quality for conservation using an integrated occurrence-mortality model. *Journal of Applied Ecology*, 46, 600-609.
- Fortin, J. K., Schwartz, C. C., Gunther, K. A., Teisberg, J. E., Haroldson, M. A., Evans, M. A., & Robbins, C. T. (2013). Dietary adjustability of grizzly bears and American black bears in Yellowstone National Park. *The Journal of wildlife management*, 77(2), 270-281.
- French, S. P., French, M. G., & Knight, R. R. (1994). Grizzly bear use of army cutworm moths in the Yellowstone ecosystem. *International Conference of Bear Research & Management*, 9, 389-399.
- GAP/LANDFIRE National Terrestrial Ecosystems. (2011). <https://maps.usgs.gov/terrestrial-ecosystems-2011/>
- Geremia, C., McGarvey, L., & White P. J. (2019). Status report on the Yellowstone bison population, 2019. Yellowstone Center for Resources, Yellowstone National Park, Mammoth, Wyoming.
- Gergel, D. R., Nijssen, B., Abatzoglou, J. T., Lettenmaier, D. P., & Stumbaugh, M. R. (2017). Effects of climate change on snowpack and fire potential in the western USA. *Climatic Change*, 141(2), 287-299.
- Graham, K., Beckerman, A. P., & Thirgood, S. (2005). Human–predator–prey conflicts: ecological correlates, prey losses and patterns of management. *Biological Conservation*, 122(2), 159-171.
- Green, G. 1994. Use of spring carrion by bears in Yellowstone National Park. M.S. Thesis, University of Idaho, Moscow, Idaho.
- Hansen, A. J., & Phillips, L. B. (2015). Which tree species and biome types are most vulnerable to climate change in the US Northern Rocky Mountains?. *Forest Ecology & Management*,

338, 68-83.

- Haroldson, M. A., Schwartz, C. C., Cherry, S., & Moody, D. S. (2004). Possible effects of elk harvest on fall distribution of grizzly bears in the Greater Yellowstone Ecosystem. *The Journal of Wildlife Management*, 68(1), 129-137.
- Haroldson, M. A., & Gunther, K. A. (2013). Roadside bear viewing opportunities in Yellowstone National Park: characteristics, trends, and influence of whitebark pine. *Ursus*, 24(1), 27-41.
- Haswell, P. M., Shepherd, E. A., Stone, S. A., Purcell, B., & Hayward, M. W. (2019). Foraging theory provides a useful framework for livestock predation management. *Journal for Nature Conservation*, 49, 69-75.
- Hobson, K. A., McLellan, B. N., & Woods, J. G. (2000). Using stable carbon ( $\delta^{13}\text{C}$ ) and nitrogen ( $\delta^{15}\text{N}$ ) isotopes to infer trophic relationships among black and grizzly bears in the upper Columbia River basin, British Columbia. *Canadian Journal of Zoology*, 78(8), 1332-1339.
- Hovey, F. W., & McLellan, B. N. (1996). Estimating population growth of grizzly bears from the Flathead River drainage using computer simulations of reproduction and survival rates. *Canadian Journal of Zoology*, 74(8), 1409-1416.
- Hufkens, K., Keenan, T. F., Flanagan, L. B., Scott, R. L., Bernacchi, C. J., Joo, E., ... & Richardson, A. D. (2016). Productivity of North American grasslands is increased under future climate scenarios despite rising aridity. *Nature Climate Change*, 6(7), 710-714.
- Isaak, D. J., Wollrab, S., Horan, D., & Chandler, G. (2012). Climate change effects on stream and river temperatures across the northwest US from 1980–2009 and implications for salmonid fishes. *Climatic Change*, 113(2), 499-524.
- Jacoby, M. E., Hilderbrand, G. V., Servheen, C., Schwartz, C. C., Arthur, S. M., Hanley, T. A., ... & Michener, R. (1999). Trophic relations of brown and black bears in several western North American ecosystems. *The Journal of Wildlife Management*, 63(3), 921-929.
- Johnson, S. J., & Griffel, D. E. (1982). Sheep losses on grizzly bear range. *The Journal of Wildlife Management*, 46(3), 786-790.
- Jorgensen, C. J. (1983). Bear-sheep interactions, Targhee National Forest. *International*

- Conference on Bear Research & Management, 5, 191-200.
- Karlsson, J., & Johansson, Ö. (2010). Predictability of repeated carnivore attacks on livestock favours reactive use of mitigation measures. *Journal of Applied Ecology*, 47(1), 166-171.
- Khorozyan, I., & Waltert, M. (2019a). How long do anti-predator interventions remain effective? Patterns, thresholds and uncertainty. *Royal Society Open Science*, 6(9), 190826.
- Khorozyan, I., & Waltert, M. (2019b). A framework of most effective practices in protecting human assets from predators. *Human Dimensions of Wildlife*, 24(4), 380-394.
- Knight, R. R., & Judd, S. L. (1983). Grizzly bears that kill livestock. *International Conference on Bear Research & Management*, 5, 186-190.
- Knight, R. R., & Eberhardt, L. L. (1985). Population dynamics of Yellowstone grizzly bears. *Ecology*, 66(2), 323-334.
- Knight, R. R., Blanchard, B. M., & Eberhardt, L. L. (1988). Mortality patterns and population sinks for Yellowstone grizzly bears, 1973-1985. *Wildlife Society Bulletin*, 16(2), 121-125.
- Lamb, C. T., Mowat, G., McLellan, B. N., Nielsen, S. E., & Boutin, S. (2017). Forbidden fruit: human settlement and abundant fruit create an ecological trap for an apex omnivore. *Journal of Animal Ecology*, 86(1), 55-65.
- Lennox, R. J., Gallagher, A. J., Ritchie, E. G., & Cooke, S. J. (2018). Evaluating the efficacy of predator removal in a conflict-prone world. *Biological Conservation*, 224, 277-289.
- Littell, J. S., Elsner, M. M., Mauger, G. S., Lutz, E. R., Hamlet, A. F., & Salathé, E. P. (2011). Regional climate and hydrologic change in the northern US Rockies and Pacific Northwest: internally consistent projections of future climate for resource management. Project report for USFS JVA 09-JV-11015600-039. Climate Impacts Group, University of Washington, Seattle, Washington.
- Liu, Z., & Wimberly, M. C. (2016). Direct and indirect effects of climate change on projected future fire regimes in the western United States. *Science of the Total Environment*, 542, 65-75.
- Lubow, B. C., & Smith, B. L. (2004). Population dynamics of the Jackson elk herd. *Journal of Wildlife Management*, 68(4), 810-829.

- Luce, C. H. (2018). Effects of climate change on snowpack, glaciers, and water resources in the Northern Rockies. Pages 25-36 in J. Halofsky & D. L. Peterson (eds.). *Climate Change and Rocky Mountain Ecosystems*. Springer, Cambridge, Massachusetts.
- Lute, M. L., Carter, N. H., López-Bao, J. V., & Linnell, J. D. (2018). Conservation professionals agree on challenges to coexisting with large carnivores but not on solutions. *Biological Conservation*, 218, 223-232.
- Mace, R., & Roberts, L. (2012). Northern Continental Divide Ecosystem grizzly bear monitoring team annual report, 2012. Montana Fish, Wildlife & Parks, Kalispell, Montana.
- Macfarlane, W. W., Logan, J. A., & Kern, W. R. (2013). An innovative aerial assessment of Greater Yellowstone Ecosystem mountain pine beetle-caused whitebark pine mortality. *Ecological Applications*, 23(2), 421-437.
- MacNulty, D. R., Stahler, D. R., Wyman, C. T., Ruprecht, J., & Smith, D. W. (2016). The challenge of understanding northern Yellowstone elk dynamics after wolf reintroduction. *Yellowstone Science*, 24(1), 25-33.
- Marcus, W. A., Meacham, J. E., Rodman, A. W., & Steingisser, A. Y. (2012). *Atlas of Yellowstone*. University of California Press, Berkeley, California.
- Mattson, D. J. (1990). Human impacts on bear habitat use. *International Conference of Bear Research & Management*, 8, 33-56.
- Mattson, D. J., Blanchard, B. M., & Knight, R. R. (1991a). Food habits of Yellowstone grizzly bears, 1977–1987. *Canadian Journal of Zoology*, 69(6), 1619-1629.
- Mattson, D. J., Gillin, C. M., Benson, S. A., & Knight, R. R. (1991b). Bear feeding activity at alpine insect aggregation sites in the Yellowstone ecosystem. *Canadian Journal of Zoology*, 69(9), 2430-2435.
- Mattson, D. J., Blanchard, B. M., & Knight, R. R. (1992). Yellowstone grizzly bear mortality, human habituation, and whitebark pine seed crops. *The Journal of Wildlife Management*, 56(3), 432-442.
- Mattson, D.J. (1997). Use of ungulates by Yellowstone grizzly bears *Ursus arctos*. *Biological Conservation* 81: 161-177.

- Mattson, D. J. (1998). Changes in mortality of Yellowstone's grizzly bears. *Ursus*, 10, 129-138.
- Mattson, D.J. (2000). Causes and consequences of dietary differences among Yellowstone grizzly bears (*Ursus arctos*). Ph.D. Dissertation, University of Idaho, Moscow, ID. 173 pp.
- Mattson, D.J., & T. Merrill (2002). Extirpations of grizzly bears in the contiguous United States, 1850–2000. *Conservation Biology* 16: 1123-1136.
- Mattson, D.J., K. Barber, R. Maw & R. Renkin (2004). Coefficients of Productivity for Yellowstone's Grizzly Bear Habitat. U.S. Geological Survey, Biological Resources Discipline Biological Science Report USGS/BRD/BSR-2002-0007.
- McLellan, B. N. (2011). Implications of a high-energy and low-protein diet on the body composition, fitness, and competitive abilities of black (*Ursus americanus*) and grizzly (*Ursus arctos*) bears. *Canadian Journal of Zoology*, 89(6), 546-558.
- Merrill, T., & Mattson, D. (2003). The extent and location of habitat biophysically suitable for grizzly bears in the Yellowstone region. *Ursus*, 14(2), 171-187.
- Merrill, T. (2005). Grizzly bear conservation in the Yellowstone to Yukon region. Yellowstone to Yukon Conservation Initiative, Technical Report, 6. Canmore, Alberta, Canada.
- Middleton, A. D., Morrison, T. A., Fortin, J. K., Robbins, C. T., Proffitt, K. M., White, P. J., ... & Kauffman, M. J. (2013). Grizzly bear predation links the loss of native trout to the demography of migratory elk in Yellowstone. *Proceedings of the Royal Society of London B: Biological Sciences*, 280(1762), 20130870.
- Miller, J. R., Stoner, K. J., Cejtin, M. R., Meyer, T. K., Middleton, A. D., & Schmitz, O. J. (2016). Effectiveness of contemporary techniques for reducing livestock depredations by large carnivores. *Wildlife Society Bulletin*, 40(4), 806-815.
- Morehouse, A. T., & Boyce, M. S. (2017). Troublemaking carnivores: conflicts with humans in a diverse assemblage of large carnivores. *Ecology and Society*, 22(3).
- Moreira-Arce, D., Ugarte, C. S., Zorondo-Rodríguez, F., & Simonetti, J. A. (2018). Management tools to reduce carnivore-livestock conflicts: current gap and future challenges. *Rangeland Ecology & Management*, 71(3), 389-394.
- Mowat, G., Heard, D. C., & Schwarz, C. J. (2013). Predicting grizzly bear density in western



- North America. PLoS One, 8(12).
- Murie, A. (1948). Cattle on grizzly bear range. *The Journal of Wildlife Management*, 12(1), 57-72.
- Niedziałkowska, M., Hayward, M. W., Borowik, T., Jędrzejewski, W., & Jędrzejewska, B. (2019). A meta-analysis of ungulate predation and prey selection by the brown bear *Ursus arctos* in Eurasia. *Mammal Research*, 64(1), 1-9.
- Northern Yellowstone Cooperative Wildlife Working Group (2016-2019). Winter surveys of Northern Yellowstone elk. e.g.: <https://www.nps.gov/yell/learn/news/2019-late-winter-survey-of-northern-yellowstone-elk.htm>
- Northrup, J. M., Stenhouse, G. B., & Boyce, M. S. (2012). Agricultural lands as ecological traps for grizzly bears. *Animal Conservation*, 15(4), 369-377.
- Orozco, K. & Miles, N. (2013) Use of diminished whitebark pine resources by adult female grizzly bears in Togwotee Pass, Spread Creek, and Mount Leidy in the Bridger-Teton National Forest, Wyoming, 2012. Pages 45-47 in F.T. Van Manen, M.A. Haroldson, & K. West (eds.). *Yellowstone grizzly bear investigations: Annual report of the Interagency Grizzly Bear Study Team*, 2012. U.S. Geological Survey, Bozeman, Montana.
- Pease, C. M., & Mattson, D. J. (1999). Demography of the Yellowstone grizzly bears. *Ecology*, 80(3), 957-975.
- Prevéy, J. S., Parker, L. E., Harrington, C. A., Lamb, C. T., & Proctor, M. F. (2020). Climate change shifts in habitat suitability and phenology of huckleberry (*Vaccinium membranaceum*). *Agricultural & Forest Meteorology*, 280, 107803.
- Proulx, G. (2018). Concerns about mammal predator killing programs: scientific evidence and due diligence. *Canadian Wildlife Biology & Management*, 7, 59.
- Ray, A. M., Sepulveda, A. J., Irvine, K. M., Wilmoth, S. K., Thoma, D. P., & Patla, D. A. (2019). Wetland drying linked to variations in snowmelt runoff across Grand Teton and Yellowstone national parks. *Science of the Total Environment*, 666, 1188-1197.
- Reeves, M. C., Moreno, A. L., Bagne, K. E., & Running, S. W. (2014). Estimating climate change effects on net primary production of rangelands in the United States. *Climatic*

Change, 126(3-4), 429-442.

- Rehfeldt, G. E., Crookston, N. L., Sáenz-Romero, C., & Campbell, E. M. (2012). North American vegetation model for land - use planning in a changing climate: A solution to large classification problems. *Ecological Applications*, 22(1), 119-141.
- Robison, H. L. (2009). Relationships between army cutworm moths and grizzly bear conservation. Ph.D. Dissertation, University of Nevada-Reno, Reno, Nevada.
- Sagør, J. T., Swenson, J. E., & Røskoft, E. (1997). Compatibility of brown bear *Ursus arctos* and free-ranging sheep in Norway. *Biological Conservation*, 81(1-2), 91-95.
- Schwartz, C. C., & Haroldson, M. A., eds. (1999). *Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 1998*. U.S. Geological Survey, Bozeman, Montana.
- Schwartz, C. C., & Haroldson, M. A., eds. (2000). *Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 1999*. U.S. Geological Survey, Bozeman, Montana.
- Schwartz, C. C., & Haroldson, M. A., eds. (2001). *Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2000*. U.S. Geological Survey, Bozeman, Montana.
- Schwartz, C. C., & Haroldson, M. A., eds. (2002). *Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2001*. U.S. Geological Survey, Bozeman, Montana.
- Schwartz, C. C., & Haroldson, M. A., eds. (2003). *Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2002*. U.S. Geological Survey, Bozeman, Montana.
- Schwartz, C. C., & Haroldson, M. A., & West, K., eds. (2004). *Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2003*. U.S. Geological Survey, Bozeman, Montana.
- Schwartz, C. C., & Haroldson, M. A., & West, K., eds. (2005). *Yellowstone grizzly bear*

- investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2004. U.S. Geological Survey, Bozeman, Montana.
- Schwartz, C. C., & Haroldson, M. A., & West, K., eds. (2006). Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2005. U.S. Geological Survey, Bozeman, Montana.
- Schwartz, C. C., & Haroldson, M. A., & West, K., eds. (2007). Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2006. U.S. Geological Survey, Bozeman, Montana.
- Schwartz, C. C., & Haroldson, M. A., & West, K., eds. (2008). Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2007. U.S. Geological Survey, Bozeman, Montana.
- Schwartz, C. C., & Haroldson, M. A., & West, K., eds. (2009). Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2008. U.S. Geological Survey, Bozeman, Montana.
- Schwartz, C. C., & Haroldson, M. A., & West, K., eds. (2010). Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2009. U.S. Geological Survey, Bozeman, Montana.
- Schwartz, C. C., & Haroldson, M. A., & West, K., eds. (2011). Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2010. U.S. Geological Survey, Bozeman, Montana.
- Schwartz, C. C., Haroldson, M. A., White, G. C., Harris, R. B., Cherry, S., Keating, K. A., ... & Servheen, C. (2006). Temporal, spatial, and environmental influences on the demographics of grizzly bears in the Greater Yellowstone Ecosystem. *Wildlife Monographs*, 161(1).
- Schwartz, C. C., Haroldson, M. A., & White, G. C. (2010). Hazards affecting grizzly bear survival in the Greater Yellowstone Ecosystem. *The Journal of Wildlife Management*, 74(4), 654-667.
- Schwartz, C. C., Fortin, J. K., Teisberg, J. E., Haroldson, M. A., Servheen, C., Robbins, C. T., & Van Manen, F. T. (2013). Body and diet composition of sympatric black and grizzly bears in

- the Greater Yellowstone Ecosystem. *Journal of Wildlife Management*, 78(1), 68-78.
- Storer, T. I., & Tevis, L. P., Jr. (1996). *California grizzly*. University of California Press, Berkeley, California.
- Swan, G. J., Redpath, S. M., Bearhop, S., & McDonald, R. A. (2017). Ecology of problem individuals and the efficacy of selective wildlife management. *Trends in Ecology & Evolution*, 32(7), 518-530.
- Thoma, D. P., Munson, S. M., Rodman, A. W., Renkin, R., Anderson, H. M., & Wacker, S. D. (2019). Patterns of primary productivity and ecological drought in Yellowstone. *Yellowstone Science*, 27(1), <https://www.nps.gov/articles/patterns-of-primary-production-ecological-drought-in-yellowstone.htm>
- Treves, A., & Naughton-Treves, L. (2005). Evaluating lethal control in the management of human-wildlife conflict. Pages 86-106 in R. Woodruffe, S. Thirgood, & A. Robinowitz (eds.). *People and Wildlife, Conflict or Coexistence?* Cambridge University Press, Cambridge, United Kingdom.
- Treves, A., Krofel, M., & McManus, J. (2016). Predator control should not be a shot in the dark. *Frontiers in Ecology & the Environment*, 14(7), 380-388.
- Van Manen, F. T., Haroldson, M. A., & West, K., eds. (2012). *Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2011*. U.S. Geological Survey, Bozeman, Montana.
- Van Manen, F. T., Haroldson, M. A., & West, K., eds. (2013). *Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2012*. U.S. Geological Survey, Bozeman, Montana.
- Van Manen, F. T., Haroldson, M. A., West, K., & Soileau, S. C., eds. (2014). *Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2013*. U.S. Geological Survey, Bozeman, Montana.
- Van Manen, F. T., Haroldson, M. A., & Soileau, S. C., eds. (2015). *Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2014*. U.S. Geological Survey, Bozeman, Montana.


- Van Manen, F. T., Haroldson, M. A., & Karabensh, B. E., eds. (2016). Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2015. U.S. Geological Survey, Bozeman, Montana.
- Van Manen, F. T., Haroldson, M. A., & Karabensh, B. E., eds. (2017). Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2016. U.S. Geological Survey, Bozeman, Montana.
- Van Manen, F. T., Haroldson, M. A., & Karabensh, B. E., eds. (2018). Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2017. U.S. Geological Survey, Bozeman, Montana.
- Van Manen, F. T., Haroldson, M. A., & Karabensh, B. E., eds. (2019). Yellowstone grizzly bear investigations: Annual Report of the Interagency Grizzly Bear Study Team: 2018. U.S. Geological Survey, Bozeman, Montana.
- Van Manen, F. T., Haroldson, M. A., Bjornlie, D. D., Ebinger, M. R., Thompson, D. J., Costello, C. M., & White, G. C. (2016). Density dependence, whitebark pine, and vital rates of grizzly bears. *Journal of Wildlife Management*, 80, 300-313.
- Vulla, E., Hobson, K. A., Korsten, M., Leht, M., Martin, A. J., Lind, A., ... & Saarma, U. (2009, December). Carnivory is positively correlated with latitude among omnivorous mammals: evidence from brown bears, badgers and pine martens. *Annales Zoologici Fennici*, 46(6), 395-415.
- Wells, S. L. (2017). Livestock depredation by grizzly bears on Forest Service grazing allotments in the Greater Yellowstone Ecosystem. M.S. Thesis, Montana State University, Bozeman, Montana.
- Wells, S. L., McNew, L. B., Tyers, D. B., Van Manen, F. T., & Thompson, D. J. (2019). Grizzly bear depredation on grazing allotments in the Yellowstone Ecosystem. *The Journal of Wildlife Management*, 83(3), 556-566.
- Westerling, A. L., Turner, M. G., Smithwick, E. A., Romme, W. H., & Ryan, M. G. (2011). Continued warming could transform Greater Yellowstone fire regimes by mid-21st century. *Proceedings of the National Academy of Sciences*, 108(32), 13165-13170.

- White, D., Jr., Kendall, K. C., & Picton, H. D. (1998) Grizzly bear feeding activity at alpine army cutworm moth aggregation sites in northwest Montana. *Canadian Journal of Zoology*, 76, 221-227.
- White, P. J., Wallen, R. L., & Hallac, D. E. (2015). Yellowstone bison: conserving an American icon in modern society. Yellowstone Association, Yellowstone National Park, Wyoming.
- Williams, J. E., Haak, A. L., Neville, H. M., & Colyer, W. T. (2009). Potential consequences of climate change to persistence of cutthroat trout populations. *North American Journal of Fisheries Management*, 29(3), 533-548.
- Wilson, S. M., Madel, M. J., Mattson, D. J., Graham, J. M., Burchfield, J. A., & Belsky, J. M. (2005). Natural landscape features, human-related attractants, and conflict hotspots: a spatial analysis of human–grizzly bear conflicts. *Ursus*, 16(1), 117-129.
- Wilson, S. M., Madel, M. J., Mattson, D. J., Graham, J. M., & Merrill, T. (2006). Landscape conditions predisposing grizzly bears to conflicts on private agricultural lands in the western USA. *Biological Conservation*, 130(1), 47-59.
- Wyoming Game & Fish Department, Annual Big Game Job Completion Reports.  
<https://wgfd.wyo.gov/Hunting/Job-Completion-Reports>
- Wyoming Game & Fish Department (2016). Wyoming grizzly bear management plan. Wyoming Game & Fish Department, Laramie, Wyoming.
- Zager, P., & Beecham, J. (2006). The role of American black bears and brown bears as predators on ungulates in North America. *Ursus*, 17(2), 95-108.


**Attachment 2. Figures Referenced in Main Text.**

**[Begin on following page]**























**Figure 5.** Trends in numbers of reproductive grizzly bear females and livestock-related conflicts in the Greater Yellowstone Ecosystem (GYE): (a) estimated numbers of females with cubs-of-the-year (COY) ecosystem-wide based on the Chao2 method as a 3-year moving average (green line) and as individual annual estimates (green squares) together with total number of livestock-related human-grizzly bear conflicts, including conflicts involving cattle, sheep, horses swine, and other smaller domestic animals (Schwartz & Haroldson 1999-2004; Schwartz et al. 2005-2011; Van Manen et al. 2012-2019); and (b) estimated numbers of females with COY relative to total numbers of cattle- and sheep-related conflicts in the GYE (burgundy line and dots) as well as conflicts only within the Upper Green River area (red line and dots) (Schwartz et al. 2008-2011; Van Manen et al. 2012-2019). Note the rapid escalation of livestock-related conflicts beginning around 2009 and the lack of corresponding increase in grizzly bear population trend.





**Figure 6.** Trends in estimated numbers of reproductive grizzly bear females, ecosystem-wide, by two different methods (Mark-resight and Chao2) relative to trend in total known and probable deaths of adolescent and adult grizzly bear in the Greater Yellowstone Ecosystem expressed as a three-year running average (Schwartz & Haroldson 2003-2004; Schwartz et al. 2005-2011; Van Manen et al. 2012-2019). Estimated numbers of females with cubs-of-the-year, as a three-year running average or three-year sum, are used to monitor trend of the Yellowstone grizzly bear population (Van Manen et al., 2019). Note the only slight increase in numbers of reproductive females at the same time that numbers of known and probable grizzly bear deaths increased substantially. By deduction, if population size increased only slightly at the same time that deaths escalated, grizzly bear death rates must have increased, especially after 2009, suggesting that mortality was becoming increasingly unsustainable.

